

AMINO CROP

Organic plant-based fertilizer containing amino acids

Product identification

Trade name **AMINOCROP**
Type of product **EC fertilizer.**

Composition (w / w%)

Total organic matter75.00%
Free amino acids..... 30.00%
Total nitrogen (N)15.00%

Properties

AMINO CROP is a completely natural and organic product. It contains 17 free amino acid types in "L" form. Recommended to stimulate the growth and development of any type of plant

Doses and applications

CROPS	Moments of application	Quantity (Kg / Ha)	Number of applications
Sugar Beet, Sunflower, Salads, Parsley	Stage 3 to 5 Leaves and 3 Weeks After	0.5-1	2 applications
Cotton, Fraise, Houblon	Stade 3 à 5 Feuilles, Croissance Végétative, Floraison	0.5-0.6-0.7	3 applications
Mais	Plants at 30 cm in height, Before the formation of the tillers	0.5-0.7	2 applications
Tobacco, Tomato,	Stage 3 to 5 Leaves, Early bloom, After: Every 4 Weeks	0.5-0.6-0.7	3 applications
Potato, Vegetables, Soybean, Canola, Fruit Tree with Stone	Before bloom, 3 weeks after.	0.5-0.7	2 applications
Grain	Stage 3 to 5 Leaves, With herbicide	0.5	2 applications
Melon, watermelon, cucumber	Stage 3 to 5 Leaves, Flowering Period, Fruit Magnification	0.5-0.6-0.7	3 applications
Citrus, Apple - Pear, Cauliflower, Broccoli	Flowering, Maturity of the fruits, 3 weeks after	0.5,-0.6	3 applications
Table grapes, Artichoke	Shoots of 30 cm, before flowering, after flowering	0.5	3 applications
Olivier	Bourgeons at 10 cm, Before 3 Weeks of harvest	0.5-0.6	2 applications
banana tree	Start Growth, 3 weeks after	0.5-0.6	2 applications
Eggplant, Chili	Stage 3 to 5 Leaves, Flowering Period, Fruiting, After: Every 3 weeks	0.5-0.6-0.7	3 applications
Onion, Carrot, Radish	2-3 weeks after germination, every 2 weeks	0.4-0.5	2 applications
Grass	Early May, mid-July, mid-September	0.4-0.5-0.6	3 applications
CROPS UNDER GREENHOUSE			
Application	foliar	fertigation	period

First Application	600 g / Ha	200g / 100lt of water	Planting
Second Application	600-800 g / Ha	200g / 100lt of water	2 weeks after
After	600 g / Ha	200g / 100lt of water	Every 10 days

Applications must be done early in the morning or evening.
Avoid applications during high rainfall and windy days.
It is best to apply the product at low and frequent doses.
Product applied via the different irrigation systems.

Safety instructions

Keep out of reach of children

Do not eat and drink during application

Do not breathe product vapor

Keep away from food, beverages and feed

In case of contact with eyes, rinse immediately with plenty of water and seek medical advice

Wear protective gloves and eye / face protection.

In case of accident or if you feel unwell, seek medical advice immediately (show the label where possible)